

10. Se sospetto una encefalopatia che quadro elettroencefalografico mi aspetto di trovare?

In Word come si "aggiunge un collegamento ipertestuale"?

One akinetic event per patient was included; if multiple akinetic events occurred in their history or during a video-EEG recording, we considered the first recorded event. Exclusion criteria included 1) a suspected or confirmed diagnosis of epilepsy; and 2) dissociative events lasting less than 20 s, ensuring that the minimum EEG duration was consistent with that analyzed in previous studies.



11. Esporre un montaggio SEP 4 canali per lo studio dell'arto superiore.

In Word come si "inserisce un sommario ad un documento"?

The video-EEGs were recorded using Xltech, employing a 21-channel 10–20 montage. The reference electrode was placed at Cz. Impedances were maintained below 5 k Ω . Electrophysiological data were continuously recorded with a sampling rate of 1000 Hz and a bandwidth of 0.05–100 Hz. For patients recorded at Policlinico Umberto I (12 patients), the recordings were part of a 30-minute routine EEG, while at Neuromed (10 patients), they were extracted from in-hospital sessions of prolonged monitoring (24 h) for diagnostic purposes.



3. Esistono metodiche di comparazione motoria per la valutazione di un possibile intrappolamento nervoso al polso di mediano e/o ulnare? Se sì quali?

In Excel come si "inserisce un grafico"?

Given the heterogeneity of this condition in clinical presentation and triggering factors, numerous psychopathological theories have emerged over the decades.

Currently, the most accepted theory is the Integrated Cognitive Model (ICM), which suggests that an emotional trigger, along with heightened autonomic arousal, leads to a loss of inhibitory control and the activation of automatic motor or behavioral responses, resulting in non-epileptic seizures.

